

TARGETS IN HETEROCYCLIC SYSTEMS

Chemistry and Properties

Volume 12 (2008)

Table of Contents

Synthetic accessibility to protoporphyrin-IX, chlorophyll *a* and (2*R*)-phytychromobilin and other biologically important pyrrole derivatives in any ¹³C and ¹⁵N enriched isotopic form

1

Prativa B. S. Dawadi and Johan Lugtenburg

1. Introduction

2. Synthesis and discussion

2.1. Isotopically labeled pyrroles, protoporphyrins and related derivatives

2.1.1. [1-¹⁵N]-3-Cyano-4-methyl-1*H*-pyrrole

2.1.2. Benzyl 3-(2-methoxycarbonylethyl)-4-methoxycarbonylmethyl-5-methyl [2-¹³C]-pyrrole-2-carboxylate, benzyl 3-(2-methoxycarbonylethyl)-4-methoxycarbonylmethyl[2-¹³C]-pyrrole-2-carboxylate, benzyl 5-(¹³C)-**formyl**)-3-(2-methoxycarbonyl-ethyl)-4-methoxy-carbonylmethyl[2-¹³C]-pyrrole-2-carboxylate and 2-(¹³C)-**formyl**)-4-(2-methoxy-carbonylethyl)-3-methoxycarbonylmethyl[5-¹³C]-pyrrole

2.1.3. Ethyl 3,5-dimethyl-4-ethyl[1-¹⁵N]-pyrrole-2-carboxylate and ethyl 3,5-dimethyl-4-ethyl[3-¹³C]-pyrrole-2-carboxylate

2.1.4. [11-¹³C]-Porphobilinogen

2.1.5. Dibenzyl 3,5-dimethyl[2,3,4,5-¹³C₄]-pyrrole-2,5-dicarboxylate

2.1.6. 5-Substituted 2-pyrrole esters

2.1.7. 3-Cyanopyrrole, 3-cyano-4-methylpyrrole, pyrrole-3-carboxylate and 4-methylpyrrole-carboxylate

2.1.8. [2-¹³C]-3-Cyano-4-methyl-3-pyrrolin-2-one and [3-¹³C]-3-cyano-4-methyl-3-pyrrolin-2-one

2.1.9. 3,4-Disubstituted pyrrolinones and 3-(**pentadeuteroethyl**)-4-methylpyrrolin-2-one

2.1.10. Chlorophyll *a*

2.1.11. Protoporphyrin-IX, hemin and biliverdin

2.1.12. (2*R*)-Phytychromobilin and (2*R/S*)-phytycyanobilin

2.1.12.1. The synthesis of pyrromethenone **124** (A,B-ring segment of phytychromobilin)

2.1.12.2. The synthesis of pyrromethenone **132** (C,D-ring segment of phytychromobilin)

2.1.12.3. Coupling of pyrromethenones **124** and **132** to form (2*R*)-phytychromobilin

2.1.13. Bilirubin and its analogs

3. Conclusion

Acknowledgments

References

□-Alkylidene-□-lactones and lactams via radical- and transition metal-mediated cyclizations

31

Laurence Feray and Michèle P. Bertrand

1. Introduction

2. Radical cyclizations

2.1. Intramolecular addition of alkoxy carbonyl radicals to conjugated alkyne

2.2. Enyne cyclization

2.2.1. Tandem addition of nucleophilic radicals to activated triple bond/5-exo ring closure

2.2.2. Tandem addition of electrophilic radicals to double bond/5-exo ring closure onto conjugated alkyne

2.3. Iodine atom transfer-mediated cyclization onto conjugated alkyne

2.4. Two-step strategies involving 5-exo-dig cyclizations followed by oxidation of resulting 3-alkylidene-tetrahydrofurans

3. Transition metal-mediated cyclizations

3.1. Cyclization of alkoxy carbonyl- and carbamoyl- metal complexes onto triple bond

3.2. Enyne cyclization

3.2.1. Palladium-catalyzed reactions

3.2.1.1. Halopalladation

3.2.1.2. Acetoxypalladation

3.2.1.3. Carbopalladation

3.2.1.4. Tandem cyclization/Suzuki coupling

3.2.2. Rhodium(I)-catalyzed reactions

3.2.2.1. Cycloisomerization of 1,6-enynes

3.2.2.2. Reductive cycloisomerization of 1,6-enynes and acetylenic aldehydes

3.2.3. Alkenylative cyclization promoted by ruthenium catalysts

3.3. Two-step procedures involving Pd(0)-mediated cyclization followed by oxidation

3.4. Miscellaneous methodologies

3.4.1. Cycloalkenylation of tungsten-alkynyl amines

3.4.2. Cyclization involving Pd(0) as the leaving group

3.4.2.1. Pd-catalyzed heteroannulation of 1,3-dienes

3.4.2.2. Pd-catalyzed carbonylation of 3-iodohomoallylic alcohols

3.4.3. Ti(II)-mediated cyclization of homopropargylic carbonates

3.4.4. [2+2+1] Cycloaddition leading to bicyclic □-alkylidene-□-lactones and lactams

4. Conclusion

The reaction of carbonyl compounds and nitriles with triflic anhydride.

59

Synthesis of pyrimidines, benzothiazines, oxazoles and triazines

Roberto Martínez-Álvarez and Antonio Herrera

1. Introduction

1.1. Triflic anhydride and vinyl triflates

2. Initial investigations

2.1. The reaction of vinyl triflates and nitriles. Mechanism of the reaction

3. Synthesis of substituted pyrimidines

3.1. The reaction of aldehydes and ketones in the presence of triflic anhydride

3.2. Regioselectivity in the reaction of aliphatic ketones

3.3. Synthesis of halo- and alkoxy pyrimidines

3.4. Synthesis of substituted benzoquinazolines and related compounds

3.5. Cyclobutapyrimidines and pyrimidine *ortho*-quinodimethanes in the Diels-Alder reaction

3.6. The special reactivity of benzyl cyanides

3.7. The use of methyl thiocyanate. Synthesis of functionalized pyrimidines

3.8. Synthesis of heterocycles fused pyrimidines

4. The reaction with carbonyl compounds containing sulfur atoms

4.1. The reaction of thioesters with nitriles and triflic anhydride

4.2. Stereoselective synthesis of substituted 1,3-benzothiazines

4.3. Synthesis of 1,3-oxazoles

5. Synthesis of 2,4,6-trisubstituted 1,3,5-triazines

6. Conclusions

Acknowledgments

References

Metal-mediated C–C and C–N bond formation in the synthesis of bioactive purines

85

Lise-Lotte Gundersen

1. Introduction

2. C-C Bond formation

2.1. Addition of organometallic reagents followed by rearomatization

2.1.1. Addition of organolithium reagents

2.1.2. Addition of organomagnesium reagents

2.2. Cross-coupling of halopurines

2.2.1. Coupling with organolithium reagents

2.2.2. Coupling with organocuprates-copper-mediated couplings

2.2.3. Coupling with organomagnesium reagents

2.2.4. Coupling with organotin reagents (Stille coupling)

2.2.5. Coupling with organozinc reagents (Negishi coupling)

2.2.6. Coupling with organoboron reagents (Suzuki-Miyaura coupling)

- 2.2.7. Coupling with organoaluminium reagents
- 2.2.8. Heck coupling
- 2.2.9. Sonogashira coupling
- 2.2.10. Direct CH alkylation and arylation
- 2.2.11. Regioselectivity in cross coupling reactions of dihalopurines
- 2.2.12. Further transformation of coupling products
- 2.3. C-C Bond formation *via* metallated purines
 - 2.3.1. Generation and reactivity of lithiated purines
 - 2.3.2. Generation and reactivity of stannylated purines
 - 2.3.3. Generation and reactivity of zincated purines
 - 2.3.4. Generation and reactivity of magnesiated purines
 - 2.3.5. Generation and reactivity of palladated purines
- 3. C-N Bond formation
 - 3.1. Buchwald-Hartwig coupling
 - 3.1.1. Buchwald-Hartwig reaction on halopurines
 - 3.1.2. Buchwald-Hartwig reaction on aminopurines
 - 3.2. Cu-mediated *N*-arylation
 - 3.3. Palladium(0)-mediated *N*-allylation
- 4. Biological activities
 - 4.1. Cytotoxic purines
 - 4.2. Antimicrobial purines
 - 4.3. Antiviral purines
 - 4.4. Adenosine receptor ligands
 - 4.5. Cytokinin analogs
 - 4.6. Inhibitors of 15-lipoxygenase
- 5. Conclusions
- Acknowledgments
- References
- Abbreviations

Indole phytoalexins from brassicaceae: synthesis and anticancer activity

120

Peter Kutschy and Roman Mezencev

- 1. Introduction
- 2. Structure and synthesis
 - 2.1. Phytoalexins with sulfur-containing side chain or substituent at indole nucleus
 - 2.2. Phytoalexins with a side chain or substituent at indole nucleus not containing sulfur
 - 2.3. 3-(Thiazole-2-yl)indole derived phytoalexins
 - 2.4. 2,3-Fused indole phytoalexins
 - 2.5. Spiroindoline[3,5']- and spiroindoline[2,5']dihydrothiazole type phytoalexins
- 3. Anticancer activity

- 3.1. Cytostatic and cytotoxic effects
- 3.2. Cancer chemopreventive activity
- 3.3. Modulation of immune response

4. Conclusions

Acknowledgment

References

Synthesis of phosphepines and application in asymmetric catalysis

149

Natalia V. Dubrovina, Ivan A. Shuklov and Armin Börner

1. Introduction

2. General strategies for the synthesis of phosphepines

2.1. Cyclization of a single component

2.1.1. Thermal cyclization of arenylphosphonic acid

2.1.2. Electrophilic cyclization

2.1.3. Reductive coupling *via* McMurry reaction

2.1.4. *Via* ring-closing metathesis

2.2. Ring formation by cyclization of two components

2.2.1. *Via* reaction of 1,6-dilithium salts with dichlorophosphines

2.2.2. *Via* reaction of silylphosphines with acyl chloride

2.2.3. *Via* radical cyclization reactions of silylphosphines with alkenes

2.2.4. Nucleophilic addition reaction between 1,5-hexadiene and phosphine

2.3. Ring-expansion

2.3.1. From phospholes and dihydrophospholes

2.3.2. From phosphinine derivatives *via* ring enlargement by a carbene unit

2.4. Synthesis and modification of chiral dinaphthophosphepines

2.4.1. Monodentate phosphepines

2.4.2. Bidentate phosphepines

3. Asymmetric catalysis with phosphepines

3.1. Asymmetric catalysis with transition metal complexes

3.1.1. Asymmetric hydrogenation

3.1.1.1. Monodentate phosphepines

3.1.1.2. Bidentate phosphepines

3.1.1.3. Asymmetric hydrogenation with self-assembling phosphepine catalyst

3.1.2. Rh-catalyzed asymmetric hydroformylation

3.1.3. Pd-catalyzed asymmetric allylic alkylation and amination (Tsuji-Trost reaction)

3.1.3.1. Pd-catalyzed asymmetric allylic alkylation

3.1.3.2. Pd-catalyzed asymmetric allylic amination

3.1.4. Enantioselective catalytic allylation of carbonyl groups by umpolung of \square -allyl palladium complexes

3.1.5. Pd-mediated Suzuki-Miyaura coupling

3.1.6. Asymmetric hydroboration

- 3.1.7. Rh-catalyzed 1,4-additions of arylboronic acids to enones
- 3.1.8. Pt-catalyzed alkoxy- and hydroxycyclization of enynes
- 3.2. Organocatalysis
 - 3.2.1. Asymmetric acylations and benzoylations
 - 3.2.2. Asymmetric [3+2] cycloadditions of imines with allenes
 - 3.2.3. Asymmetric [4+2] cycloadditions of imines with allenes

References and notes

Cyclic phospho di- and triester as structural elements of nucleic acids

185

Christelle Dupouy, Corinne Payrastré and Jean-Marc Escudier

1. Introduction
 - 1.1. Nucleotides conformation
 - 1.2. Nucleic acids primary and secondary structures
 - 1.3. Modified nucleic acids
 2. Macrocyclic phosphate esters
 - 2.1. U-Turn model
 - 2.2. 5'-Cyclouridylic acid and derivatives
 - 2.3. Synthesis of cyclic di- and trinucleotide by Ring Closure Metathesis (RCM)
 3. Dioxaphosphorinane-Constrained Nucleic Acid
 - 3.1. Diastereoselective synthesis of □□-D-CNA
 - 3.2. Synthesis of □□□-D-CNA and □□□-D-CNA
 - 3.3. Nucleosidic and sugar based approach towards □□-D-CNA
 - 3.4. Structural assignment
 - 3.5. Preliminary results on DNA duplex stabilization
 4. Conclusions
- Acknowledgments
- References

Chemistry of biologically active □-carbolines

212

Marta Rosillo, Alvaro González-Gómez, Gema Domínguez and Javier Pérez-Castells

1. Introduction
 - 1.1. Naturally occurring □-carbolines
 - 1.2. Biological activity of □-carbolines
2. Synthesis of □-carbolines
 - 2.1. The Pictet-Spengler reaction for the construction of the pyridine ring
 - 2.1.1. Catalytic PS reaction
 - 2.1.2. Solid-phase supported PS
 - 2.1.3. Pictet-Spengler combined with other processes
 - 2.2. The Bischler-Napieralski reaction

- 2.3. Organometallic based methods
- 2.4. Construction of the pyrrole ring
- 2.5. Cycloaddition methodologies
- 2.6. Other strategies
- 3. Transformations of β -carbolines
 - 3.1. Oxidations
 - 3.2. Reductions
 - 3.3. Stereoselective alkylations
 - 3.4. Metathesis
 - 3.5. Reactions with activated alkynes
 - 3.6. β -Carbolines as chiral ligands
- 4. Total syntheses of natural alkaloids containing the β -carboline system
- 5. Conclusions
- Acknowledgments
- References

Meso-substituted porphyrin synthesis from monopyrrole: an overview

258

Mariette M. Pereira, Carlos J. P. Monteiro and Andreia F. Peixoto

- 1. Introduction
- 2. *Meso*-substituted porphyrin synthesis
 - 2.1. One-pot synthesis
 - 2.2. Two-step synthesis
- 3. *Meso*-tetraarylporphyrin synthesis using sustainable chemistry approach
 - 3.1. Solvent-free reactions
 - 3.2. Microwave synthesis
 - 3.3. Ionic liquids
 - 3.4. Heterogeneous catalysis: zeolites and clays

Acknowledgments

References

Abbreviations

1,2- And 1,3-oxazolopyridines: versatile building blocks for the heterocyclic synthesis

279

Gianluca Giorgi and Fabio Ponticelli

- 1. Introduction
 - 1.1. Availability
 - 1.1.1. Oxazolopyridines
 - 1.1.2. Isoxazolopyridines
- 2. Structural characterization
 - 2.1. Nuclear Magnetic Resonance
 - 2.2. X-Ray crystallography

- 2.3. Mass spectrometry
- 3. Chemical properties
 - 3.1. General aspects
 - 3.2. Oxazolopyridines reactivity
 - 3.3. Isoxazolopyridine reactivity
 - 3.3.1. Thermal reactivity
 - 3.3.2. Photochemical reactions
- 4. Conclusions
- Acknowledgments
- References

**New methods for the asymmetric synthesis of piperidines and pyrrolidines:
chiral auxiliaries and asymmetric organocatalysis**

302

Jose L. Vicario, Dolores Badía, Luisa Carrillo, Nerea Ruiz and Efraim Reyes

- 1. Introduction
- 2. Asymmetric synthesis of piperidine alkaloids using (*S,S*)-(+)-pseudoephedrine as chiral auxiliary
 - 2.1. A general and stereodivergent method for the asymmetric synthesis of 2-alkylpiperidine alkaloids
 - 2.2. Asymmetric synthesis of (–)-β-*conhydrine*
- 3. Organocatalytic asymmetric synthesis of pyrrolidines
 - 3.1. Organocatalytic enantioselective [3+2] cycloaddition of azomethine ylides and α,β-unsaturated aldehydes
 - 3.2. Organocatalytic Michael reaction of aldehydes and nitroacrolein dimethylacetal. Synthesis of 3,4-disubstituted and 2,3,4-trisubstituted pyrrolidines
- 4. Concluding remarks
- Acknowledgments
- References

Synthesis of phosphorus-substituted sulfur heterocycles

328

Mihaela Gulea and Annie-Claude Gaumont

- 1. Introduction
- 2. Syntheses by phosphorylation of a sulfur heterocycle
 - 2.1. Syntheses using phosphorus nucleophiles
 - 2.1.1. Arbuzov-type reactions
 - 2.1.2. Metal-catalyzed P–C coupling reactions
 - 2.1.3. Nucleophilic additions of di- or trialkylphosphites
 - 2.1.4. Pummerer-type reactions
 - 2.2. Syntheses using phosphorus electrophiles
- 3. Syntheses involving the formation of the sulfur heterocycle

- 3.1. Cycloaddition reactions
 - 3.1.1. [4+2] Cycloadditions
 - 3.1.1.1. With phosphorylated thiocarbonyl compounds as dienophiles
 - 3.1.1.1.1. Phosphinoyl thioaldehydes
 - 3.1.1.1.2. Phosphonodithioformates
 - 3.1.1.1.3. Phosphoryl-substituted sulfines
 - 3.1.1.2. With phosphorylated thiocarbonyl compounds as dienes
 - 3.1.2. [3+2] Cycloadditions
 - 3.2. Syntheses based on intramolecular cyclizations
 - 3.2.1. Carbanion mediated cyclizations
 - 3.2.2. Carbenoid mediated cyclizations
 - 3.2.3. Radical mediated cyclizations
 - 3.2.4. Electrocyclizations
 - 3.2.4.1. 1,3-Dipolar electrocyclizations
 - 3.2.4.2. Domino Knoevenagel/1,6-heteroelectrocyclizations
 - 3.3. Miscellaneous
 - 3.3.1. Sigmatropic rearrangements
 - 3.3.1.1. [2,3]-Sigmatropic rearrangement
 - 3.3.1.2. [1,3]-Sigmatropic rearrangement
 - 3.3.2. Ring-closing metathesis
4. Conclusion
- Acknowledgments
- References

Carbolithiation reactions in the synthesis of heterocycles

349

Roberto Sanz and Alberto Martínez

- 1. Introduction
- 2. Intermolecular reactions
 - 2.1. General aspects of intermolecular carbolithiation reactions
 - 2.2. Synthesis of indole derivatives
 - 2.3. Synthesis of quinoline derivatives
 - 2.4. Synthesis of other heterocyclic systems
- 3. Intramolecular reactions
 - 3.1. Special features of intramolecular carbolithiation reactions
 - 3.1.1. Stereoselectivity and mechanism
 - 3.1.2. Scope and limitations
 - 3.2. Synthesis of furan derivatives
 - 3.3. Synthesis of benzofuran derivatives
 - 3.4. Synthesis of pyrrole derivatives
 - 3.5. Synthesis of azabicyclo derivatives

- 3.6. Synthesis of indole derivatives
- 3.7. Synthesis of other heterocyclic systems
- 3.8. Synthesis of heterocycles through carbolithiation of arynes

4. Conclusions

Acknowledgments

References

Oligo- and poly(2,5-thienylene-ethynylene)s

390

Herbert Meier

- 1. Introduction
- 2. Synthetic approach
 - 2.1. Monodisperse oligomers
 - 2.2. Multidisperse polymers
- 3. Molecular structures
- 4. Electronic absorption and fluorescence
- 5. Applications in materials science
- 6. Conclusion and outlook

References and notes

Chemoenzymatic synthesis of optically active pyridine derivatives

414

Vicente Gotor-Fernández, Eduardo Busto and Vicente Gotor

- 1. Introduction
- 2. Use of hydrolases
 - 2.1. Lipase-catalyzed kinetic and dynamic kinetic resolution of alcohols or esters
 - 2.2. Proteases in the resolution of esters through hydrolytic processes
 - 2.3. Asymmetrization or desymmetrization of diols or diacetates using lipases
 - 2.4. Kinetic resolution of racemic amines using lipases in organic solvents
 - 2.5. Classical kinetic resolutions using epoxide hydrolases
- 3. Use of oxidoreductases
 - 3.1. Alcohol dehydrogenases
 - 3.1.1. Bioreductions
 - 3.1.2. Biooxidations
 - 3.1.3. Deracemizations combining biooxidation and bioreduction processes
 - 3.2. Bioxygenations using oxygenases
- 4. Use of lyases
 - 4.1. Synthesis of chiral cyanohydrins using oxynitrilases
 - 4.2. Aldolases for the production of chiral pyridines

5. Concluding remarks

Acknowledgments

References

Asymmetric synthesis of substituted piperazines

438

Jean-Charles Quirion

1. Introduction
 2. Resolution and diastereomeric separation
 - 2.1. Chemical resolution
 - 2.2. Enzymatic resolution
 - 2.3. Diastereomeric separation
 3. Reduction of 2-keto and diketopiperazines
 - 3.1. Reduction of 2-ketopiperazines
 - 3.2. Reduction of 2,3-diketopiperazines
 - 3.3. Reduction of 2,5-diketopiperazines
 4. Reduction of tetrahydropyrazines
 - 4.1. Reduction of chiral tetrahydropyrazines
 - 4.2. Asymmetric hydrogenation of tetrahydropyrazines
 5. Reductive cyclization of diamines
 - 5.1. Diastereoselective cyclization
 - 5.2. Enantioselective cyclization
 6. Cyclization of diaminoalcohols
 7. Cyclization of diamines
 - 7.1. Nucleophilic substitution
 - 7.2. Reductive amination
 - 7.3. Allylic substitution
 8. Amination and carboamination of olefins
 9. Direct functionalization of substituted piperazines
 - 9.1. Selective alkylation of substituted piperazines
 - 9.2. *N*-Selective functionalization
 10. Conclusions
- References

Microwave assisted synthesis of functional oligothiophenes

460

Manuela Melucci and Giovanna Barbarella

1. Introduction
 - 1.1. Oligothiophenes scenario
 - 1.2. Ring-by ring growth through metal catalyzed cross-coupling reaction
2. Solvent-free microwave assisted Suzuki coupling
 - 2.1. Unsubstituted quater- and quinquethiophene
 - 2.2. Self-affinity and electrical properties of **T5**
 - 2.3. Methyl substituted sexithiophene
3. Liquid phase microwave assisted Suzuki coupling (less soluble oligomers)

- 3.1. Rigid core thiophene co-oligomers with liquid crystalline properties
 - 4. Microwave assisted synthesis of oligothiophene semiconductors in aqueous media using silica and chitosan supported Pd catalysts
 - 4.1. Optimization of reaction conditions
 - 4.2. Extension to other substrates
 - 5. Conclusions
- References

ISBN 978-88-86208-56-7 ISSN 1724-9449

XV+485 pp

Euro 80.00 + postal expenses