

TARGETS IN HETEROCYCLIC SYSTEMS

Chemistry and Properties

Volume 6 (2002)

Table of Contents

Enantioselective ring expansion via aziridinium intermediates. Synthesis of substituted piperidines from substituted pyrrolidines. Synthetic applications

Janine Cossy and Domingo Gomez Pardo

1. Introduction
 2. Mechanism
 3. Synthesis of 3-halogenopiperidines
 4. Synthesis of 3-oxygenated and 3-amino-piperidines derivatives
 5. Synthesis of C(3)-alkylated piperidine derivatives
 6. Conclusions
- Acknowledgements
References

The formation of the carbon-carbon bond catalysed by metalloporphyrins

Pietro Tagliatesta

1. Introduction
 - 1.1. Catalysis by metalloporphyrins: general remarks
 2. Cyclopropanation reactions
 - 2.1. Rhodium porphyrin catalysts and asymmetric cyclopropanation reactions
 - 2.2. Iron porphyrin catalysts
 - 2.3. Osmium porphyrin catalysts
 - 2.4. Ruthenium porphyrin catalysts and asymmetric cyclopropanation reactions
 3. Diels-Alder reaction
 - 3.1. Catalysis by aluminium porphyrins
 4. Olefinations of aldehydes
 - 4.1. Catalysis by iron porphyrins
 5. Cyclotrimerization of acetylene derivatives
 - 5.1. Substituted benzenes from acetylenes: catalysis by iron porphyrins
 - 5.2. Substituted benzenes from acetylenes: catalysis by rhodium porphyrins
 6. Conclusions
- Acknowledgements
References

Pyrazoles as drugs: facts and fantasies

José Elguero, Pilar Goya, Nadine Jagerovic and Artur M. S. Silva

1. Introduction
2. Agents acting on the central nervous system
 - 2.1. Antidepressants, antipsychotics, anxiolytics, sedatives and hypnotics
 - 2.1.1. Antidepressants
 - 2.1.2. Antipsychotics
 - 2.1.3. Anxiolytics
 - 2.1.4. Sedatives and hypnotics
 - 2.2. Anticonvulsants, analgesics, anti-Parkinson and anti-Alzheimer drugs
 - 2.2.1. Anticonvulsants
 - 2.2.2. Analgesics
 - 2.2.3. Anti-Parkinson drugs
 - 2.2.4. Anti-Alzheimer drugs
 - 2.3. Pyrazoles acting on the cannabinoid receptors
3. Pharmacodynamic agents

- 3.1. Antiarrhythmics, antianginals, vasodilators, antihypertensives, diuretics, antithrombotics and anti-allergics
 - 3.1.1. Antiarrhythmics, antianginals, vasodilators, antihypertensives
 - 3.1.2. Diuretics
 - 3.1.3. Antithrombotics
 - 3.1.4. Anti-allergics
 - 3.2. Drugs acting on gastrointestinal, respiratory and urogenital systems
 - 3.2.1. Antihistaminics
 - 3.2.2. Antiemetics
 - 3.2.3. Respiratory system
 - 3.2.4. Urogenital system
 - 3.3. Drugs acting on skin diseases
 - 3.3.1. Psoriasis
 4. Agents acting on metabolic diseases and on endocrine functions
 - 4.1. Anti-inflammatory drugs and antiarthritics
 - 4.2. Hypoglycemic, hypolipidemic and antiobesity agents
 - 4.3. Peptide and steroidal hormones
 - 4.4. Liver alcohol dehydrogenase inhibitors
 5. Chemotherapeutic agents
 - 5.1. Anticancer drugs
 - 5.2. Immunosuppressants and immunostimulants
 - 5.3. Antiviral, antibacterial, antiparasitic, antiprotozoa and fungicides
 - 5.3.1. Antiviral
 - 5.3.2. Antibacterial
 - 5.3.3. Antiparasitic and antiprotozoa
 - 5.3.4. Fungicides
 6. Conclusions: structure-biological activity relationship and modelling
- Acknowledgements
- References

Recent advances in the synthesis of saturated nitrogen heterocycles using *N*-acyliminium ion intermediates

Marino Petrini

1. Introduction
 2. *N*-Acyliminium ions
 - 2.1. General aspects
 - 2.2. Precursors of *N*-acyliminium ions
 - 2.3. Acidic catalysts and nucleophiles
 3. Synthesis of saturated nitrogen heterocycles
 - 3.1. Pyrrolidines
 - 3.2. Pyrrolidinones
 - 3.3. Piperidines
 - 3.4. Indolizidines
 - 3.5. Quinolizidines
 - 3.6. Other bicyclic systems
 - 3.7. Tricyclic systems
 - 3.8. Polycyclic systems
 4. Conclusions
- Acknowledgements
- References

Reductive opening of heterocycles with lithium metal as a source of functionalised organolithium compounds: synthetic applications

Miguel Yus and Francisco Foubelo

1. Introduction
 2. Reductive opening of strained heterocycles
 - 2.1. Reductive opening of epoxides
 - 2.2. Reductive opening of aziridines
 - 2.3. Reductive opening of oxetanes
 - 2.4. Reductive opening of azetidines
 3. Reductive opening of cyclic allyl and benzyl ethers, amines and thioethers
 - 3.1. Reductive opening of cyclic allyl ethers, amines and related thioethers
 - 3.2. Reductive opening of cyclic benzyl ethers
 - 3.3. Reductive opening of cyclic benzyl amines
 - 3.4. Reductive opening of cyclic benzyl thioethers
 4. Reductive opening of cyclic aryl ethers and thioethers
 - 4.1. Reductive opening of cyclic aryl ethers
 - 4.2. Reductive opening of cyclic aryl thioethers
 6. Conclusions
- References

Furan-2,3-diones: convenient synthons in heterocyclic chemistry

Gert Kollenz and Sabine Leber

1. Introduction
2. Synthesis and preparation
3. Thermolysis (pyrolysis)
 - 3.1. Formation of neat α -oxoketenes
 - 3.2. *In situ*-generation of α -oxoketenes
 - 3.3. α -Oxoketenes-chemical reactions
 - 3.3.1. Dimerization
 - 3.3.2. Electrocyclization
 - 3.3.3. Cycloadditions
 - 3.3.4. Reactions with nucleophiles
4. Cycloaddition Reactions
 - 4.1. 5-Aryl-furan-2,3-diones
 - 4.2. 4-Acyl-furan-2,3-diones
 - 4.2.1. Thermal cycloaddition reactions
 - 4.2.2. Photocycloaddition reactions
5. Reactions with nucleophiles
 - 5.1. 5-Aryl-furan-2,3-diones
 - 5.1.1. NH-nucleophiles
 - 5.1.2. OH-nucleophiles
 - 5.1.3. C-nucleophiles
 - 5.2. 4-Acyl-furan-2,3-diones
 - 5.2.1. NH₂-nucleophiles
 - 5.2.2. Miscellaneous

References

Metal-catalyzed cyclocarbonylation reactions for the synthesis of lactones and lactams

Giuseppe Vasapollo and Giuseppe Mele

1. Introduction
2. Cyclocarbonylation reactions
 - 2.1. General features on cyclocarbonylation reactions
 - 2.2. Synthesis of lactones
 - 2.2.1. Synthesis of four-membered lactones (β -lactones)
 - 2.2.2. Synthesis of five-membered lactones
 - 2.2.2.1. Synthesis of five-membered unsaturated lactones (furanones)

- 2.2.2.2. Synthesis of five- and six-membered saturated lactones
- 2.2.3. Synthesis of five-, six- and seven membered fused to aromatic rings lactones and bis-lactones
- 2.3. Synthesis of lactams and pyrrolidinones
 - 2.3.1. Synthesis of four-membered lactams (β -lactams)
 - 2.3.2. Synthesis of five-membered lactams
 - 2.3.2.1. Synthesis of five-membered unsaturated lactams
 - 2.3.2.2. Synthesis of five-membered saturated lactams (pyrrolidinones)
 - 2.3.3. Synthesis of six-membered unsaturated lactams
 - 2.3.4. Synthesis of five- and six-membered fused to aromatic rings lactams
- 2.4. The Pauson-Khand reaction
- 3. Trends in cyclocarbonylation reactions
 - 3.1. Cyclocarbonylation of steroids and other natural substrates
 - 3.2. More atoms in the heterocyclic system
 - 3.3. 8,9-Membered and larger rings
 - 3.4. Heterogeneous catalyzed cyclocarbonylation reactions
 - 3.5. Enantioselective cyclocarbonylation reactions
 - 3.6. Cyclocarbonylation products as building blocks for new materials
- 4. Conclusions
- References

Preparation and synthetic applications of (*S*)- and (*R*)-*N*-BOC-*N,O*-isopropylidene- α -methylserinals

Alberto Avenzoa, Jesús H. Busto, Carlos Cativiela, Francisco Corzana, Jesús M. Peregrina, David Sucunza and María M. Zurbano

- 1. Introduction
- 2. Synthesis of (*S*)- and (*R*)-*N*-Boc-*N,O*-isopropylidene- α -methylserinals
- 3. Synthetic applications
 - 3.1. α -Methyl- α -amino acids
 - 3.1.1. Enantiomerically pure α -substituted alanines
 - 3.1.1.1. (*R*)- and (*S*)-Isovaline
 - 3.1.1.2. Unsaturated α -substituted alanines: vinylalanines and ethynylalanines
 - 3.1.2. α -Methyl- α -amino acids with two stereogenic centres
 - 3.2. Carbohydrates of glycopeptide antibiotics
- 4. Conclusions
- Acknowledgements
- References

Synthesis and reactivity of 1-(2,4,6-trialkylphenyl)phospholes with a flattened P-pyramid

György Keglevich

- 1. Introduction
 - 1.1. General considerations
 - 1.2. Synthesis of phospholes
 - 1.3. Aromaticity of phospholes
- 2. Synthesis of 1-(2,4,6-trialkylphenyl)phospholes
- 3. Aromaticity of 1-(2,4,6-trialkylphenyl)phospholes
- 4. Reactivity of 1-(2,4,6-trialkylphenyl)phospholes
 - 4.1. Aromatic electrophilic substitutions
 - 4.2. Reactions with phosphorustribromide
 - 4.3. Diels-Alder cycloadditions
 - 4.4. Sigmatropic rearrangements
 - 4.5. Complexation reactions
 - 4.6. Oxidation reactions
- 5. Phosphole ligands in rhodium complexes

6. Fragmentation of phosphole-based phosphanorbornenes
 7. Conclusions
- Acknowledgement
References

Chiral β -aminoalcohols and derivatives in the asymmetric synthesis of the tetrahydroisoquinolines

Eneritz Anakabe, Dolores Badía, Luisa Carrillo, Efraim Reyes and Jose L. Vicario

1. Introduction
 - 1.1. β -Aminoalcohols and derivatives as chiral building blocks
 - 1.2. β -Aminoalcohols and derivatives as chiral auxiliaries
 - 1.3. β -Aminoalcohols and derivatives as chiral ligands
 2. (*S*)-Phenylglycinol as chirality source
 - 2.1. Asymmetric synthesis of 3-aryltetrahydroisoquinolines
 - 2.2. Diastereodivergent synthesis of 1,3-disubstituted tetrahydroisoquinolines
 - 2.3. Asymmetric synthesis of protoberberines
 - 2.4. Asymmetric synthesis of isopavines
 - 2.5. Asymmetric synthesis of 4-alkyltetrahydroisoquinolines
 3. (*S*)-Arylglycinols as chiral templates
 - 3.1. Synthesis of 3-aryl-4-hydroxytetrahydroisoquinolines
 - 3.2. General procedure for the asymmetric synthesis of 3-aryltetrahydroisoquinolines
 4. (*S,S*)-(+)-Pseudoephedrine as chiral auxiliary
 - 4.1. Asymmetric synthesis of 4-alkyl-3-aryl-1,2,3,4-tetrahydroisoquinolines
 - 4.2. Asymmetric synthesis of B/C-hexahydrobenzo[*c*]phenanthridines
 5. Concluding remarks
- Acknowledgements
References

Stereoselective synthesis of nitrogen heterocycles of enantiopure 3-hydroxy esters through electrophilic amination

Christine Greck and Olivia Poupardin

1. Introduction
 2. Diastereoselective synthesis of *anti* 2-hydrazino-3-hydroxy esters
 - 2.1. Electrophilic amination of 1,3-dioxan-4-ones
 - 2.2. Electrophilic amination of 3-hydroxy esters
 3. Synthesis of nitrogen heterocycles
 - 3.1. Six-membered rings
 - 3.1.1. (3*S*, 4*S*)-4-Hydroxy-2,3,4,5-tetrahydropyridazine-3-carboxylic acid
 - 3.1.2. *Trans* 3-hydroxypipercolic acid and application to (-)-swainsonine
 - 3.2. Five-membered rings
 - 3.2.1. *Trans* 3-hydroxy-*D*-proline
 - 3.2.2. (4*S*, 5*R*)-5-Carbomethoxy-4-hydroxy- Δ^2 -pyrazoline
 - 3.3. Four-membered rings
 - 3.3.1. *Cis* 4-methyl-3-AMA
 - 3.3.2. *Cis* 4-carbamoyloxymethyl-3-AMA
 4. Conclusion
- References

Recent trends in synthesis of heterocycles using ketene-dithioacetals with electron attracting groups

Galal H. Elgemeie and Ghada W. Abd-Elaziz

1. Introduction
2. Synthesis
3. Utility in heterocyclic synthesis
 - 3.1. Five-membered rings

- 3.1.1. Five-membered rings with one hetero-atom
- 3.1.2. Five-membered rings with two hetero-atoms
- 3.1.3. Five-membered rings with three hetero-atoms
- 3.2. Six-membered rings
 - 3.2.1. Six-membered rings with one hetero-atom
 - 3.2.2. Six-membered rings with two hetero-atoms
 - 3.2.3. Six-membered rings with three hetero-atoms
- 3.3. Seven-membered rings
- 3.4. Fused heterocyclic compounds

References

Recent development in the chemistry of 4,5-dihalo-3(2H)-pyridazinones

Pál Tapolcsányi and Péter Mátyus

- 1. Introduction
- 2. Substitution at the lactam moiety
 - 2.1. Alkylated derivatives
 - 2.2. Sulfonylation
 - 2.3. Acylation
- 3. Halogen exchange reactions
 - 3.1. Nucleophilic substitution reactions
 - 3.2. Palladium catalysed C-C coupling reactions
- 4. Concluding remarks

Acknowledgements

References

Synthesis of heterocycles from allenes by intramolecular ring closure

Jacques Hatem

- 1. Introduction
- 2. Heterocycles with three-membered rings
 - 2.1. Epoxydation
 - 2.2. Aziridination
- 3. Heterocycles with four-membered rings
- 4. Heterocycles with five-membered rings
 - 4.1. Oxygen heterocycles
 - 4.1.1. Saturated oxygen heterocycles
 - 4.1.2. Unsaturated oxygen heterocycles
 - 4.1.2.1. Dihydrofurans
 - 4.1.2.2. Furans
 - 4.1.2.3. Furanones
 - 4.2. Nitrogen heterocycles
 - 4.2.1. Saturated nitrogen heterocycles
 - 4.2.2. Unsaturated nitrogen heterocycles
 - 4.3. Other heterocycles
- 5. Heterocycles with six-membered rings
 - 5.1. Oxygen heterocycles
 - 5.2. Nitrogen heterocycles
- 6. Heterocycles with larger rings
- 7. Miscellaneous
- 8. Conclusion

References