

Table of Contents

Preparation of biomedically interesting heterocycles starting from oxiranic compounds	1
<i>Florenci V. González, Lledó Bou-Iserte</i>	
1. Introduction	
2. Synthesis of nitrogen heterocycles starting from nitroepoxides	
3. Synthesis of thiazoles, piperidones and quinoxalines starting from cyanoepoxides	
4. Synthesis of morpholines starting from epoxides	
5. Synthesis of quinoxalines, tetrahydroquinoxalines and pyrazines starting from epoxides	
6. Synthesis of azetidines starting from epoxides	
7. Synthesis of piperidines and pyrrolidines from azidoepoxides	
8. Synthesis of heterocycles starting from epoxyesters and epoxyketones	
9. Other reactions involving synthesis of heterocycles from oxiranic compounds	
10. Conclusion	
Acknowledgement	
References	
Photocatalyzed preparation of oxygenated heterocycles	17
<i>Stefano Crespi, Stefano Protti, Paolo Quadrelli, Maurizio Fagnoni, Davide Ravelli</i>	
1. Introduction	
2. Three- and four-membered rings	
3. Five-membered rings	
3.1. Tetrahydrofurans and dihydrobenzofurans	
3.2. γ -Lactones and butenolides	
3.3. Furans and benzofurans	
4. Six-membered rings	
4.1. Chroman derivatives	
4.2. δ -Lactones and coumarins	
4.3. Other oxygen heterocycles	
5. Larger than six-membered rings	
6. Conclusions	
Acknowledgements	
References	
Recent advances in the PdI₂-catalyzed carbonylative synthesis of heterocycles from acetylenic substrates: a personal account	41
<i>Bartolo Gabriele</i>	
1. Introduction	
2. PdI ₂ -catalyzed oxidative heterocyclization-alkoxycarbonylation	
3. PdI ₂ -catalyzed oxidative aminocarbonylation-heterocyclization	
4. PdI ₂ -catalyzed non-oxidative carbonylations	
5. Conclusion	
Acknowledgement	
References	
Recent advances in metal-catalyzed C-H functionalization of pyrimidinones, quinazolinones and fused quinazolinones	56
<i>Corinne Fruit, Thierry Besson</i>	
1. Introduction	
2. Metal-catalyzed C-H functionalization of pyrimidinones	
2.1. Intramolecular palladium-catalyzed C-H arylation of protected uracils using aryl halides	
2.2. Intermolecular metal-catalyzed C-H arylation of protected uracils	

- 2.3. Metal-catalyzed dehydrogenative cross-coupling of pyrimidinones with (hetero)-arenes
 - 2.4. Palladium-catalyzed dehydrogenative alkenylation of uracils
 - 2.5. Miscellaneous functionalization of uracils and pyrimidinone
 - 3. Metal-catalyzed C-H functionalization of quinazolinones
 - 4. Metal-catalyzed C-H functionalization of 2-arylquinazolinones
 - 5. Metal-catalyzed C-H functionalization of fused quinazolinones
 - 6. Conclusion
- Acknowledgments
Abbreviations
References

The chemistry of non-S-oxidised 4*H*-1,2,6-thiadiazines

82

Andreas S. Kalogirou, Panayiotis A. Koutentis

- 1. Introduction
- 2. Theoretical methods
- 3. Experimental structural methods
 - 3.1. X-Ray diffraction
 - 3.2. NMR spectroscopy
 - 3.3. UV/vis spectroscopy
 - 3.4. Infrared and Raman spectroscopy
 - 3.5. Mass spectrometry
- 4. Thermodynamic aspects
 - 4.1. Physical properties
 - 4.2. Aromaticity
- 5. 1,2,6-Thiadiazine ring synthesis
 - 5.1. Synthesis of monocyclic 4*H*-1,2,6-thiadiazines
 - 5.2.1. Synthesis of polycyclic 4*H*-1,2,6-thiadiazines by building of the thiadiazine ring
 - 5.2.2. Synthesis of polycyclic 4*H*-1,2,6-thiadiazines by fusion onto the thiadiazine ring
- 6. Chemistry of 1,2,6-thiadiazines
 - 6.1. Functional group interconversions at the C4 position of monocyclic 4*H*-1,2,6-thiadiazines
 - 6.1.1. Interconversions of 3,4,4,5-tetrachloro-4*H*-1,2,6-thiadiazine **18**
 - 6.1.1.1. Conversion to 2-(3,5-dichloro-4*H*-1,2,6-thiadiazin-4-ylidene)malononitrile **4**
 - 6.1.1.2. Conversion to 4*H*-1,2,6-thiadiazin-4-imines
 - 6.1.1.3. Conversion to 3,5-dichloro-4*H*-1,2,6-thiadiazin-4-one **5**
 - 6.1.1.4. Conversion to ketals and thioketals
 - 6.1.1.5. Other conversions
 - 6.1.2. Interconversions of 4*H*-1,2,6-thiadiazin-4-ones
 - 6.1.3. Interconversions of 2-(4*H*-1,2,6-thiadiazin-4-ylidene)malononitriles **61a** and **69a**
 - 6.1.4. Interconversions of 4*H*-1,2,6-thiadiazin-4-imines
 - 6.1.5. Interconversions of 1,2,6-thiadiazine-4-ketals
 - 6.1.6. Interconversions of 4*H*-1,2,6-thiadiazin-4-thiones
 - 6.2. Functional group interconversions at the C4 position of polycyclic 4*H*-1,2,6-thiadiazines
 - 6.3. Displacement of C3/5 chlorides by nucleophiles on monocyclic 4*H*-1,2,6-thiadiazines
 - 6.3.1. Chloride displacements of 4*H*-1,2,6-thiadiazin-4-ones
 - 6.3.2. Chloride displacements of 4*H*-1,2,6-thiadiazin-4-imines
 - 6.3.3. Chloride displacements of 2-(4*H*-1,2,6-thiadiazin-4-ylidene)malononitriles
 - 6.3.4. Chloride displacements of thiadiazine-4-ketals
 - 6.4. Displacement of C3/5 chlorides by nucleophiles on polycyclic 4*H*-1,2,6-thiadiazines
 - 6.5. Pd C-C coupling of chloro-4*H*-1,2,6-thiadiazines
 - 6.5.1. Couplings of 4*H*-1,2,6-thiadiazin-4-ones
 - 6.5.1.1. Suzuki-Miyaura couplings (symmetrical)
 - 6.5.1.2. Suzuki-Miyaura couplings (unsymmetrical)

- 6.5.1.3. Sonogashira couplings
- 6.5.1.4. Homocoupling
- 6.5.1.5. Stille couplings (symmetrical)
- 6.5.1.6. Stille couplings (unsymmetrical)
- 6.5.2. Couplings of 4*H*-1,2,6-thiadiazin-4-imines
- 6.5.3. Couplings of 4*H*-1,2,6-thiadiazine-4-ketals
- 6.6. Pd-Catalysed C-N coupling of chloro-4*H*-1,2,6-thiadiazines
- 6.7. Oxidation of the 4*H*-1,2,6-thiadiazine sulfur atom
- 7. Transformations of 1,2,6-thiadiazine to other ring systems
 - 7.1. Transformations to 4,5,6-trichloropyrimidine-2-carbonitrile **121**
 - 7.2. Transformations to 1,2,5-thiadiazoles **123**
 - 7.3. Transformations to other ring systems
- 8. Degradation of the 1,2,6-thiadiazine ring
- 9. Important compounds and applications
 - 9.1. Medicinal applications
 - 9.2. Agrochemical applications
 - 9.3. Electronic applications
 - 9.4. Other applications
- 10. Conclusions
- Acknowledgements
- References

Nitroalkenes in diverse synthesis of heterocyclic compounds with two or three heteroatoms: recent advances 119

Seyyed Emad Hooshman, Azim Ziyaei Halimehjani

- 1. Introduction
- 2. Imidazoles and their reduced derivatives
- 3. Pyrazoles and pyrazolidines
- 4. Triazoles
- 5. Thiazoles and thiazines
- 6. Isoxazolines and oxazines
- 7. Dihydropyrimidines
- 8. Imidazopyridines
- 9. Bicyclic compounds
- 10. Diazepines
- 11. Reactions of nitroepoxides
- 12. Conclusion
- Acknowledgements
- References

Palladium-catalyzed domino carbopalladation/cyclization of allenes 138

Roberto Sala, Gianluigi Broggin

- 1. Introduction
- 2. Carboamination of allenes
 - 2.1. Palladium-catalyzed intermolecular coupling/intramolecular amination processes
 - 2.2. Palladium-catalyzed intramolecular coupling/intramolecular amination processes
 - 2.3. Carbonylative carbopalladation/intramolecular amination processes
- 3. Carboalkoxylation of allenes
 - 3.1. Palladium-catalyzed intermolecular coupling/intramolecular alkoxylation processes
 - 3.2. Carbonylative carbopalladation/intramolecular alkoxylation processes
- 4. Carboalkylation of allenes
- 5. Conclusions

References

- Enantioselective synthesis of five-membered heterocycles through [3+2]-cycloadditions with isocyano esters** 165
Gonzalo Blay, Carlos Vila, Pablo Martínez-Pardo, José R. Pedro
1. Introduction
 2. Enantioselective synthesis of 2-oxazolines from carbonyl compounds
 - 2.1. Enantioselective reactions of α -isocyano esters with aldehydes
 - 2.1.1. Gold and silver catalysis
 - 2.1.2. Platinum and palladium catalysis
 - 2.1.3. Organocatalytic reactions
 - 2.1.4. Cooperative metal/organocatalysis
 - 2.2. Enantioselective reactions of α -isocyano esters with ketones
 - 2.2.1. Enantioselective reactions with 1,2-dicarbonyl compounds
 - 2.2.2. Enantioselective reactions with unactivated ketones
 3. Enantioselective synthesis of 2-imidazolines from imines
 - 3.1. Enantioselective reactions of α -isocyano esters with aldimines
 - 3.1.1. Metal catalysis
 - 3.1.2. Organocatalytic reactions
 - 3.1.3. Cooperative metal/organocatalysis
 - 3.2. Enantioselective reactions of α -isocyano esters with ketimines
 - 3.2.1. Reaction with acyclic ketimines
 - 3.2.2. Reaction with cyclic and isatin ketimines
 4. Enantioselective synthesis of dihydropyrroles and related compounds
 - 4.1. Enantioselective reactions of α -isocyano esters with acyclic electrophilic alkenes
 - 4.2. Enantioselective synthesis of 3,3'-pyrrolidinyl spirooxindoles
 - 4.3. Desymmetrization of cyclic olefins
 - 4.4. Enantioselective synthesis of pyrroloindolines via cascade reactions
 5. Enantioselective synthesis of 1,2,4-triazolines
 6. Conclusion
- Acknowledgements
References
- Recent advances in the synthesis of new pyrazole derivatives** 194
Juan-Carlos Castillo, Jaime Portilla
1. Introduction
 2. Functionalized pyrazoles
 - 2.1. Aminopyrazoles
 - 2.2. Formylpyrazoles
 3. Fused pyrazoles
 - 3.1. Pyrazolo[1,5-*a*]pyrimidines
 - 3.2. Pyrazolo[3,4-*b*]pyridines
 - 3.3. Synthetic utilities of fused pyrazoles
 4. Conclusion
- Acknowledgement
References
- Synthetic approaches towards cyclopenta[*b*]indole scaffold** 224
Samia R. Lima, Hugo Santos, Ralph C. Gomes, Manoel T. Rodrigues Jr., Fernando Coelho
1. Introduction
 2. Metal-catalyzed approaches
 - 2.1. Palladium-catalyzed approaches

- 2.2. Gold-catalyzed approaches
 - 2.3. Rhodium-catalyzed approaches
 - 3. Brønsted acid catalyzed or promoted approaches
 - 4. Lewis acid catalyzed or promoted approaches
 - 4.1. Cyclization based on electrophilic aromatic substitutions (S_EAr)
 - 4.2. Cyclization based on Nazarov and Nazarov-type reactions
 - 4.2.1. Racemic approaches
 - 4.2.2. Enantioselective approaches
 - 5. Miscellaneous approaches
 - 6. Conclusions
- Acknowledgments
References

Quinolones for applications in medicinal chemistry: synthesis and structures

260

Pedro Horta, Alina Secrieru, Andy Coninckx, Maria L. S. Cristiano

- 1. Introduction
 - 2. Quinolones in medicinal chemistry
 - 2.1. Quinolones as antimalarial agents
 - 2.2. Quinolones as antibacterial agents
 - 2.3. Quinolones as antimycobacterial agents
 - 2.4. Quinolones as anticancer agents
 - 2.5. Quinolones with antiviral activity
 - 2.5.1. Quinolones as drugs against HIV
 - 2.5.2. Quinolones as drugs against HSV and HCV
 - 2.6. Quinolones with antifungal activity
 - 2.7. Quinolones with neurologic activity
 - 2.7.1. Quinolones as drugs for the treatment of anxiety
 - 2.7.2. Quinolones as drugs for the treatment of Alzheimer's disease
 - 2.7.3. Quinolones with antinociceptive activity
 - 2.8. Quinolones with immunomodulatory and anti-inflammatory activities
 - 2.9. Quinolones with metabolic activity
 - 2.10. Quinolones as drugs for the treatment of ischemia
 - 3. Synthesis of quinolones
 - 3.1. Type *A*
 - 3.2. Type *B*
 - 3.3. Type *C*
 - 3.4. Type *D*
 - 3.5. Type *E*
 - 3.6. Combination of multi-bond formation
 - 4. Liabilities of quinolones
 - 4.1. Toxicity and side effects
 - 4.2. Selectivity
 - 4.3. Resistance
 - 4.4. Drug-drug and drug-food interactions
 - 4.5. Solubility, synthesis, characterization and pharmacokinetic profile
 - 4.6. Synthesis inherent problems
 - 4.6.1. Structural isomerism in quinolone synthesis
 - 4.6.2. Oxo-quinoline/hydroxy-quinoline tautomerism
 - 5. Conclusions
- Acknowledgement
References

- Iso(thio)cyanate-strategy for the organocatalytic synthesis of selected heterocyclic structures** 298
Sebastian Frankowski, Aleksandra Więckowska, Lukasz Albrecht
1. Introduction
 2. Mechanistic considerations
 3. Application of iso(thio)cyanates in asymmetric organocatalysis
 - 3.1. Electron-withdrawing-group-activated aliphatic iso(thio)cyanates in organocatalytic cascades
 - 3.2. Oxindole-derived isothiocyanates in organocatalytic cascades
 - 3.3. Other isothiocyanates in organocatalytic cascades
 4. Conclusions
- Acknowledgements
References
- The chemistry of 3-nitrochromenes** 318
Robby Vroemans, Wim Dehaen
1. Introduction
 2. Synthesis of 3-nitrochromenes
 3. Reactivity of 3-nitrochromenes
 - 3.1. Oxidations
 - 3.2. Reductions
 - 3.3. Conjugate additions
 - 3.4. Cycloaddition reactions
 4. Conclusion
- Acknowledgement
References
- Chemistry and biology of 3,4-dihydropyrimidin-2(1H)-one (or thione) derivatives obtained by the Biginelli multicomponent reaction** 356
Brenno A. D. Neto, Talita de A. Fernandes, Mauro V. Correia
1. Introduction
 2. The Biginelli multicomponent reaction mechanisms
 3. Catalysis' roles and solvent effects
 4. Enantioselective versions of the Biginelli multicomponent reaction
 5. Biological activity of DHPM
 - 5.1. Anticancer activity
 - 5.2. Antimicrobial activity
 - 5.3. Anti-HIV activity
 - 5.4. Calcium channel blockers (CCBs)
 6. Concluding remarks
- Acknowledgement
References
- Synthesis of heterocyclic systems from α,β -unsaturated diazoketones** 377
João V. Santiago, Clarice A. D. Caiuby, Antonio C. B. Burtoloso
1. Introduction
 - 1.1. Synthesis of α -diazocarbonyl compounds
 - 1.2. Synthesis of α,β -unsaturated diazoketones
 2. Recent methods for the synthesis of α,β -unsaturated diazoketones
 3. Mechanism for the stereospecific synthesis of *E*- or *Z*- α,β -unsaturated diazoketones
 4. Synthetic application of α,β -unsaturated diazoketones
 - 4.1. General applications
 - 4.2. α,β -Unsaturated diazoketones as intermediates for the synthesis of heterocyclic
 - 4.3. Burtoloso's research group contribution in the synthesis of heterocyclic systems *via*

α,β -unsaturated diazoketones intermediates
 5. Conclusions
 Acknowledgment
 References

Isoxazol-5-ones: unusual heterocycles with great synthetic potential

409

Alessandra A. G. Fernandes, Amanda F. da Silva, Samuel Thurow, Celso Y. Okada Jr., Igor D. Jurberg

1. Introduction
 - 1.1. General reactivity
 2. Applications in organic synthesis
 - 2.1. Functionalization of the isoxazolone ring
 - 2.1.1. Alkylation
 - 2.1.2. Oxidation
 - 2.1.3. EDA complexes
 - 2.1.4. Annulation
 - 2.1.4.1. Cycloadditions
 - 2.1.4.2. Spirocyclizations
 - 2.2. Transforming the isoxazolone ring
 - 2.2.1. Alkynes
 - 2.2.1.1. Flash vacuum pyrolysis (FVP)
 - 2.2.1.2. Nitrosative treatment
 - 2.2.2. Catalytic methods involving metal-nitrenoid intermediates
 - 2.2.2.1. Azadienes
 - 2.2.2.2. 1-Azabicyclo[3.1.0]hex-2-enes and related heterocycles
 - 2.2.2.3. Pyridines and piperidines
 - 2.2.2.4. 2*H*-Azirines
 - 2.2.3. Reductive protocols
 - 2.2.3.1. Quinolines
 - 2.2.3.2. Pyridines
 - 2.2.4. Other metal-catalyzed transformations
 - 2.2.4.1. Isoquinolines
 - 2.2.4.2. 1,3-Oxazin-6-ones
 - 2.2.4.3. Pyrimidinediones
 - 2.2.5. Photochemical reactions
 - 2.2.5.1. Photolysis
 - 2.3. Total syntheses of natural products
 - 2.3.1. Isoxazolones as intermediates
 - 2.3.1.1. Total synthesis of (-)-Lycoramine
 - 2.3.2. Isoxazolones as synthetic targets
 - 2.3.2.1. Synthesis of Parnafungins A and C models
 - 2.3.2.2. Synthesis of isoxazolone glucosides
 3. Conclusions
- Acknowledgements
 References

Reactions of 5-formyl- and 5-acyl-3,4-dihydro-2*H*-pyrans and their annelated analogs with nucleophiles

436

Dmitry V. Osipov, Vitaly A. Osyanin, Yury N. Klimochkin

1. Introduction
2. Reactions of 5-formyl- and 5-acyl-3,4-dihydro-2*H*-pyrans with nucleophiles
 - 2.1. C-Nucleophiles
 - 2.1.1. Enolates and CH-acids

- 2.1.2. Organometallic compounds
- 2.1.3. 1,3-C,N-Binucleophiles
- 2.1.4. Miscellaneous C-nucleophiles
- 2.2. N-Nucleophiles
 - 2.2.1. Amines
 - 2.2.2. 1,2-N,N- and 1,2-N,O-Binucleophiles
 - 2.2.3. 1,3-N,N-Binucleophiles
- 2.3. Miscellaneous nucleophiles
- 3. Reactions of carbonyl-containing 4*H*-chromenes and their annelated analogs with nucleophiles
- 4. Conclusion
- Acknowledgement
- References

Post-Ugi transformation of *N*-substituted-2-alkyneamides for the construction of diverse heterocyclic scaffolds 468

Saeed Balalaie, Pegah Shakeri

1. Introduction
2. Synthesis of functionalized *N*-substituted-2-alkyneamides through Ugi four-component reaction (Ugi-4CR)
3. Post-Ugi transformation reaction based on *N*-substituted-2-alkyneamides
4. Metal-catalyzed post-Ugi transformation of *N*-substituted-2-alkyneamides
 - 4.1. Pd-catalyzed post-Ugi transformation of functionalized *N*-substituted-2-alkyneamides
 - 4.2. Six-component reaction for the stereoselective synthesis of 3-arylidene-2-oxindoles *via* sequential Ugi/Heck carbocyclization/Sonogashira/nucleophilic addition
 - 4.3. Sequential Ugi-4CR/C-H activation using (diacetoxyiodo)benzene for the synthesis of 3-(diphenylmethylidene)-2,3-dihydro-1*H*-indol-2-ones
5. Au-catalyzed post-Ugi transformation of functionalized *N*-substituted-2-alkyneamides
6. Metal-free post-Ugi transformation of functionalized *N*-substituted-2-alkyneamides to access heterocycles
 - 6.1. Post-Ugi transformation through radical formation
 - 6.2. Metal-free synthesis of fused triazolodiazepino[5,6-*b*]quinolone derivatives *via* a sequential Ugi-4CR/nucleophilic substitution/intramolecular click reaction
7. Conclusion
- Acknowledgement
- References